



 T-4
 5

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07D 513/06, A61K 31/435		A1	(11) International Publication Number: WO 97/40051 (43) International Publication Date: 30 October 1997 (30.10.97)
(21) International Application Number: PCT/JP97/01395	(22) International Filing Date: 23 April 1997 (23.04.97)	(81) Designated States: AL, AM, AU, AZ, BA, BB, BG, BR, BY, CA, CN, CU, CZ, EE, GE, HU, IL, IS, KG, KR, KZ, LC, LK, LR, LT, LV, MD, MG, MK, MN, MX, NO, NZ, PL, RO, RU, SG, SI, SK, TJ, TM, TR, TT, UA, US, UZ, VN, YU, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).	
(30) Priority Data: 8/102303 24 April 1996 (24.04.96) JP 8/330801 11 December 1996 (11.12.96) JP			
(71) Applicant (for all designated States except US): TAKEDA CHEMICAL INDUSTRIES, LTD. [JP/JP]; 1-1, Doshomachi 4-chome, Chuo-ku, Osaka-shi, Osaka 541 (JP).			Published With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.
(72) Inventors; and (75) Inventors/Applicants (for US only): TAKATANI, Muneo [JP/JP]; 6-1, Taniguchisonomachi, Ukyo-ku, Kyoto-shi, Kyoto 616 (JP), SHIBOUTA, Yumiko [JP/JP]; 30-20-601, Deguchi-cho, Suita-shi, Osaka 564 (JP), SUGIYAMA, Yasuo [JP/JP]; 7-2, Daiwahigashi 5-chome, Kawanishi-shi, Hyogo 666-01 (JP), KAWAMOTO, Tetsuji [JP/JP]; 26-2-B203, Nakakidacho, Neyagawa-shi, Osaka 572 (JP).			
(74) Agents: ASAHIKA, Tadao et al.; Osaka Plant of Takeda Chemical Industries, Ltd., 17-85, Jusohonmachi 2-chome, Yodogawa-ku, Osaka-shi, Osaka 532 (JP).			

(54) Title: FUSED IMIDAZOPYRIDINE DERIVATIVES AS ANTIHYPERLIPIDEMIC AGENTS



(57) Abstract

A novel compound of formula (I) wherein ring Q is an optionally substituted pyridine ring; one of R⁰, R¹ and R² is -Y⁰-Z⁰, and the other two groups are a hydrogen, a halogen, an optionally substituted hydroxy group, a hydrocarbon group that may be an optionally substituted hydrocarbon group or an acyl group; Y⁰ is a bond or an optionally substituted bivalent hydrocarbon group; Z⁰ is a basic group which may be bonded via oxygen, nitrogen, -CO-, -CS-, -SO₂N(R³)- (where R³ is hydrogen or an optionally substituted hydrocarbon group), or S(O)_n (wherein n is 0, 1 or 2); is a single bond or a double bond, or a salt thereof, which has an excellent LDL receptor up-regulating, blood-lipids lowering, blood-sugar lowering and diabetic complication-ameliorating activity.


FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

CLAIMS

1. A compound of the formula:



wherein ring Q is an optionally substituted pyridine ring:

one of R^0 , R^1 and R^2 is $-Y^0-Z^0$, and the other two groups are a hydrogen, a halogen, an optionally substituted hydroxy group, an optionally substituted hydrocarbon group or an acyl group;

Y^0 is a bond or an optionally substituted divalent hydrocarbon group;

Z^0 is a basic group which may be bonded via oxygen, nitrogen, $-CO-$, $-CS-$, $-SO_2N(R^3)-$ (wherein R^3 is a hydrogen or an optionally substituted hydrocarbon group), or


-S(O)_n- (wherein n is 0, 1 or 2); and

----- is a single bond or a double bond, or a salt thereof.

2. A compound of claim 1, wherein R^0 is $-Y^0-Z^0$, wherein Y^0 and Z^0 are of the same meanings as defined in claim 1.

3. A compound of claim 1, wherein z^0 is a group with a molecular weight of not greater than 1000.


4. A compound of claim 1 which is a compound of the formula:



wherein ring Q is an optionally substituted pyridine ring;

A and B independently are an optionally substituted divalent hydrocarbon group which may be bonded via $-\text{CON}(\text{R}^{4a})-$, $-\text{CO}-$ or $-\text{N}(\text{R}^{4a})-$;

X is a bond, oxygen, sulfur, $-\text{N}(\text{R}^5)\text{CO}-$, $-\text{CO}(\text{R}^5)-$, $-\text{CO}-$ or $-\text{N}(\text{R}^5)-$;

Y is a bond, $-\text{CH}=\text{CH}-$ or $-\text{CH}=\text{CH}-$ ;

Z is $-\text{CO}-$, $-\text{COO}-$, $-\text{CON}(\text{R}^3)-$, $-\text{SO}_2\text{N}(\text{R}^3)-$ or $-\text{S}(\text{O})^m-$ (wherein m is 0, 1 or 2);


R^1 and R^2 independently are a hydrogen, a halogen, an optionally substituted hydroxy group, an optionally substituted hydrocarbon group or an acyl group;

R^3 , R^4 , R^{4a} and R^5 independently are a hydrogen or an optionally substituted hydrocarbon group; or

R^3 and A, R^4 and A, R^{4a} and B, R^5 and R^4 , or R^4 and R may independently be bonded to each other to form a ring; R is an optionally substituted hydrocarbon group or an optionally substituted heterocyclic group; and

_____ is a single bond or a double bond, or a salt thereof.

5. A compound of claim 1 which is a compound of the formula:



wherein ring Q is an optionally substituted pyridine ring;

A and B independently are an optionally substituted divalent hydrocarbon group which may be bonded via -CON(R^{4a})-, -CO- or -N(R^{4a})-;

X is a bond, oxygen, sulfur, -N(R⁵)CO-, -CO(R⁵)-, -CO- or -N(R⁵)-;

Y is a bond, -CH=CH- or -CH=CH- ;

Z is -CO-, -COO-, -CON(R³)-, -SO₂N(R³)- or -S(O)_m- (wherein m is 0, 1 or 2);

R¹ and R² independently are a hydrogen, a halogen, an optionally substituted hydroxy group, an optionally substituted hydrocarbon group or an acyl group;

R³, R⁴, R^{4a} and R⁵ independently are a hydrogen or an optionally substituted hydrocarbon group; or

R³ and A, R⁴ and A, R⁴ and B, R⁴ and R⁵, or R⁴ and R independently may be bonded to each other to form a ring;

R is an optionally substituted hydrocarbon group or an optionally substituted heterocyclic group; and


— is a single bond or a double bond, or a salt thereof.

6. A compound of claim 5, wherein A and B independently are an alkylene group; X is a bond; and

R^3 and R^4 independently are a hydrogen or an optionally substituted alkyl, cycloalkyl, alkenyl, aralkyl or aryl group.

7. A compound of claim 5, wherein ring Q is an unsubstituted pyridine ring; X is a bond; Y is a bond, $\text{---}\text{C}\text{---}$ or $\text{---}\text{C}\text{---}\text{C}\text{---}$; A and B independently are a C_{1-15} alkylene group; R^1 and R^2 independently are a hydrogen; R^3 and R^4 independently are a hydrogen or a C_{1-15} alkyl, C_{3-8} cycloalkyl, C_{2-18} alkenyl, C_{7-16} aralkyl or C_{6-14} aryl group; and R is a C_{6-14} aryl group.

8. A compound of claim 1 which is a compound of the formula:



wherein ring Q is an optionally substituted pyridine ring;

ring Q_1 is an optionally substituted nitrogen-containing heterocyclic ring;

A' is a bond or an optionally substituted divalent hydrocarbon group which may be bonded via $-\text{CON}(R^{4a})-$, $-\text{CO}-$ or $-\text{N}(R^{4a})-$;


B is an optionally substituted divalent hydrocarbon group;

X is a bond, oxygen, sulfur, $-\text{N}(R^5)\text{CO}-$, $-\text{CO}(R^5)-$, $-\text{CO}-$ or $-\text{N}(R^5)-$;

Y is a bond, -CH=CH- or -CH=CH-;


R¹ and R² independently are a hydrogen, a halogen, an optionally substituted hydroxy group, an optionally substituted hydrocarbon group or an acyl group; R³, R⁴ and R⁵ independently are a hydrogen or an optionally substituted hydrocarbon group; or R³ and A¹ may be bonded to each other to form a ring; R is an optionally substituted hydrocarbon group or an optionally substituted heterocyclic group; and ---- is a single bond or a double bond, or a salt thereof.

9. A compound of claim 8, wherein ring Q is an unsubstituted pyridine ring; R¹ and R² are a hydrogen; R³ is a hydrogen or a C₁₋₁₅ alkyl, C₃₋₈ cycloalkyl, C₂₋₁₈ alkenyl, C₇₋₁₆ aralkyl or C₆₋₁₄ aryl group; A¹ is (i) a bond, (ii) a C₁₋₁₅ alkylene group which may be substituted by 1 to 3 substituents selected from the group consisting of hydroxy, oxo and phenyl, (iii) a C₂₋₁₆ alkenylene group or (iv) a phenylene group; B is (i) a C₁₋₁₅ alkylene group which may be substituted by 1 to 3 substituents selected from the group consisting of hydroxy, oxo and phenyl, (ii) a C₂₋₁₆ alkenylene group or (iii) a phenylene group; ring Q₁ is a group of the formula:



wherein A² is =C or CH; X is a bond, oxygen, sulfur or -CON(R⁵)-; R⁵ is a hydrogen or a C₁₋₁₅ alkyl group.

10. A compound of claim 1 which is a compound of the formula:



wherein ring Q is an optionally substituted pyridine ring;

A¹ is a bond or an optionally substituted divalent hydrocarbon group which may be bonded via -CON(R⁴a)-, -CO- or -N(R⁴a)-;

B is an optionally substituted divalent hydrocarbon group;

X is a bond, oxygen, sulfur, -N(R⁵)CO-, -CON(R⁵)-, -CO- or -N(R⁵)-;

R¹ is a hydrogen, a halogen, an optionally substituted hydroxy group, an optionally substituted hydrocarbon group or an acyl group;

R³, R⁴a and R⁵ independently are a hydrogen or an optionally substituted hydrocarbon group;

R is an optionally substituted hydrocarbon group or an optionally substituted heterocyclic group;

one of G¹ and G² is N, and the other is CH or N;

ring G² is an optionally substituted ring;

g is 0, 1 or 2; and

_____ is a single bond or a double bond, or a salt thereof.

11. A compound of claim 10, wherein ring Q is a pyridine ring which may be substituted by 1 to 3 substituents selected from the group consisting of nitro, hydroxy, cyano, carbamoyl, mono- or di-C₁₋₄

alkyl-carbamoyl, carboxy, C₁₋₄ alkoxy-carbonyl, sulfo, halogen, C₁₋₄ alkoxy, phenoxy, naphthoxy, benzyloxy, halophenoxy, C₁₋₄ alkylthio, mercapto, phenylthio, pyridylthio, C₁₋₄ alkylsulfinyl, phenylsulfinyl, C₁₋₄ alkylsulfonyl, phenylsulfonyl, amino, C₁₋₃ acylamino, mono- or di-C₁₋₄ alkylamino, C₁₋₄ alkyl and C₁₋₄ haloalkyl.

12. A compound of claim 10, A¹ is a bond or a C₁₋₁₅ alkylene, C₂₋₁₆ alkenylene group which may be bonded via -CON(R^{4a})-, -CO- or -N(R^{4a})-, wherein R^{4a} is of the same meaning as defined in claim 10.

13. A compound of claim 10, B is a C₁₋₁₅ alkylene or C₂₋₁₆ alkenylene group.

14. A compound of claim 10, X is a bond, oxygen, sulfur, -CONH- or -CO-.

15. A compound of claim 10, R¹ is (1) a hydrogen, (2) a halogen, (3) a hydroxy group which may be substituted by a C₁₋₆ alkyl, phenyl, C₇₋₁₀ aralkyl, formyl, C₁₋₆ alkyl-carbonyl, phenoxy carbonyl, C₇₋₁₀ aralkyloxy-carbonyl, pyranyl, furanyl or silyl group, (4) a C₁₋₁₅ alkyl, C₃₋₈ cycloalkyl, C₂₋₁₈ alkenyl, C₇₋₁₆ aralkyl or C₆₋₁₄ aryl group or (5) a C₁₋₆ alkoxy-carbonyl, mono-C₁₋₆ alkyl-carbamoyl, di-C₁₋₆ alkyl-carbamoyl or C₁₋₁₀ alkanoyl group.

16. A compound of claim 10, R³ is a hydrogen or a C₁₋₁₅ alkyl, C₃₋₈ cycloalkyl, C₂₋₁₈ alkenyl, C₇₋₁₆ aralkyl or C₆₋₁₄ aryl group.

17. A compound of claim 10, R is (1) a C₁₋₁₅ alkyl, C₃₋₈

cycloalkyl or C₂₋₁₈ alkenyl group which may be substituted by 1 to 5 substituents selected from the group consisting of (i) nitro, (ii) hydroxy, (iii) cyano, (iv) carbamoyl, (v) mono- or di-C₁₋₄ alkyl-carbamoyl, (vi) carboxy, (vii) C₁₋₄ alkoxy-carbonyl, (viii) sulfo, (ix) halogen, (x) C₁₋₄ alkoxy, (xi) phenoxy, (xii) halophenoxy, (xiii) C₁₋₄ alkylthio, (xiv) mercapto, (xv) phenylthio, (xvi) pyridylthio, (xvii) C₁₋₄ alkylsulfinyl, (xviii) C₁₋₄ alkylsulfonyl, (xix) amino, (xx) C₁₋₃ alkanoylamino, (xxi) mono- or di-C₁₋₄ alkylamino, (xxii) 4- to 6-membered cyclic amino, (xxiii) C₁₋₃ alkanoyl, (xxiv) benzoyl and (xxv) 5- to 10-membered heterocyclic group;

(2) a C₇₋₁₆ aralkyl group which may be substituted by 1 to 4 substituents selected from the group consisting of (i) halogen, (ii) C₁₋₄ alkyl, (iii) C₂₋₆ alkenyl, (iv) C₁₋₄ alkanoyl, (v) C₁₋₄ alkoxy, (vi) nitro, (vii) cyano, (viii) hydroxy, (ix) C₁₋₄ alkoxy-carbonyl, (x) carbamoyl, (xi) mono- or di-C₁₋₄ alkyl-carbamoyl and (xii) mono- or di-C₂₋₄ alkenyl-carbamoyl;

(3) a C₆₋₁₄ aryl group which may be substituted by 1 to 4 substituents selected from the group consisting of (i) halogen, (ii) C₁₋₄ alkyl, (iii) C₁₋₄ haloalkyl, (iv) C₁₋₄ haloalkoxy, (v) C₁₋₄ alkoxy, (vi) C₁₋₄ alkylthio, (vii) hydroxy, (viii) carboxy, (ix) cyano, (x) nitro, (xi) amino, (xii) mono- or di-C₁₋₄ alkylamino, (xiii) formyl, (xiv) mercapto, (xv) C₁₋₄ alkyl-carbonyl, (xvi) C₁₋₄ alkoxy-carbonyl, (xvii) sulfo, (xviii) C₁₋₄ alkylsulfonyl, (xix) carbamoyl, (xx) mono- or di-C₁₋₄ alkyl-carbamoyl, (xxi) oxo and (xxii) thioxo; or

(4) a 5- or 6-membered monocyclic heterocyclic group containing 1 to 4 hetero-atoms selected from oxygen, sulfur and nitrogen or a fused bicyclic heterocyclic group containing 1 to 6 hetero-atoms selected from oxygen, sulfur and nitrogen, each of which may be

substituted by 1 to 4 substituents selected from the group consisting of (i) halogen, (ii) C₁₋₄ alkyl, (iii) C₁₋₄ haloalkyl, (iv) C₁₋₄ haloalkoxy, (v) C₁₋₄ alkoxy, (vi) C₁₋₄ alkylthio, (vii) hydroxy, (viii) carboxy, (ix) cyano, (x) nitro, (xi) amino, (xii) mono- or di-C₁₋₄ alkylamino, (xiii) formyl, (xiv) mercapto, (xv) C₁₋₄ alkyl-carbonyl, (xvi) C₁₋₄ alkoxy-carbonyl, (xvii) sulfo, (xviii) C₁₋₄ alkylsulfonyl, (xix) carbamoyl, (xx) mono- or di-C₁₋₄ alkyl-carbamoyl, (xxi) oxo and (xxii) thioxo.

18. A compound of claim 10, wherein ring G^a is a ring which may be substituted by 1 or 2 substituents selected from the group consisting of oxo and C₁₋₆ alkyl.

19. A compound of claim 10, wherein ring Q is an unsubstituted pyridine ring; R¹ and R³ are a hydrogen; G¹ is CH; G² is N; g is 1; and R is an optionally substituted hydrocarbon group or an optionally substituted heterocyclic group.

20. A compound of claim 19, wherein ring G^a is unsubstituted ring.

21. A compound of claim 19, wherein A¹ is a bond or a C₁₋₆ alkylene group.

22. A compound of claim 19, wherein A¹ is a bond.

23. A compound of claim 19, wherein B is a C₁₋₆ alkylene group.


24. A compound of claim 19, X is a bond.

25. A compound of claim 10, ring Q is an unsubstituted pyridine ring; R¹ and R³ are a hydrogen; A¹ is a bond; G¹ is CH; G² is N; ring G³ is a ring which may be substituted by 1 or 2 substituents selected from the group consisting of oxo and C₁₋₆ alkyl; q is 1; B is a C₁₋₆ alkylene group; X is a bond; and R is an optionally substituted phenyl group.

26. A compound of claim 25, ring G³ is unsubstituted ring.

27. A compound of claim 25, wherein R is a phenyl group which may be substituted by 1 to 3 substituents selected from the group consisting of halogen, hydroxy, C₁₋₄ alkyl, C₁₋₄ haloalkyl, C₁₋₄ alkoxy and C₁₋₄ haloalkoxy.

28. A compound of claim 1 which is a compound of the formula:




wherein ring Q is an optionally substituted pyridine ring;
 ring Q₂ is an optionally substituted nitrogen-containing heterocyclic ring;
 A is an optionally substituted divalent hydrocarbon group which may be bounded via -CON(R^{4a})-, -CO- or -N(R^{4a})-;

Y is a bond, -CH=CH- or -CH=CH-;


R¹ and R² independently are a hydrogen, a halogen, an optionally substituted hydroxy group, an optionally substituted hydrocarbon group or an acyl group; R³ and R⁴ independently are a hydrogen or an optionally substituted hydrocarbon group; or R³ and A may be bonded to each other to form a ring; R is an optionally substituted hydrocarbon group or an optionally substituted heterocyclic group; and _____ is a single bond or a double bond, or a salt thereof.

29. A compound of claim 28, wherein ring Q is an unsubstituted pyridine ring; R¹ and R² are a hydrogen; R³ is a hydrogen or a C₁₋₁₅ alkyl, C₃₋₈ cycloalkyl, C₂₋₁₈ alkenyl, C₇₋₁₆ aralkyl or C₆₋₁₄ aryl group; A is (i) a C₁₋₁₅ alkylene group which may be substituted by 1 to 3 substituents selected from the group consisting of hydroxy, oxo and phenyl, (ii) a C₂₋₁₆ alkenylene group or (iii) a phenylene group; ring Q₂ is a group of the formula:



wherein B¹ is =C, CH or N.

30. A compound of claim 1 which is a compound of the formula:



wherein ring Q is an optionally substituted pyridine ring;

ring Q₃ and Q₄ independently are an optionally substituted nitrogen-containing heterocyclic ring; A³ and B² independently are a bond or an optionally substituted divalent hydrocarbon group; X is a bond, oxygen, sulfur, -N(R⁵)CO-, -CON(R⁵)-, -CO- or -N(R⁵)-;

Y is a bond, -CH=CH- or -CH=CH- ;

R¹ and R² independently are a hydrogen, a halogen, an optionally substituted hydroxy group, an optionally substituted hydrocarbon group or an acyl group;

R⁵ is a hydrogen or an optionally substituted hydrocarbon group;


R is an optionally substituted hydrocarbon group or an optionally substituted heterocyclic group; and

_____ is a single bond or a double bond, or a salt thereof.

31. A compound of claim 30, wherein ring Q is an unsubstituted pyridine ring; R¹ and R² are a hydrogen; A³ and B² independently are a bond or a C₁₋₁₅ alkylene, C₂₋₁₆ alkenylene or phenylene group; ring Q₃ is a group of the formula:




wherein A⁴ is =C or CH; ring Q₄ is a group of the formula:



wherein A⁵ is =C or CH.

32. A compound of claim 1 which is a compound of the formula:



wherein ring Q is an optionally substituted pyridine ring;

ring Q₅ is an optionally substituted nitrogen-containing heterocyclic ring;

A is an optionally substituted divalent hydrocarbon group which may be bonded via -CON(R^{4a})-, -CO- or -N(R^{4a})-;


Y is a bond, -CH=CH- or -CH=CH- ;

R¹ and R² independently are a hydrogen, a halogen, an optionally substituted hydroxy group, an optionally substituted hydrocarbon group or an acyl group;

R³ and R^{4a} independently are a hydrogen or an optionally substituted hydrocarbon group; and

_____ is a single bond or a double bond, or a salt thereof.

33. A compound of claim 32, wherein ring Q is an unsubstituted pyridine ring; R¹ and R² are a hydrogen; R³ is a hydrogen or a C₁₋₁₅ alkyl, C₃₋₈ cycloalkyl, C₂₋₁₈ alkenyl, C₇₋₁₆ aralkyl or C₆₋₁₄ aryl group; A is a C₁₋₁₅ alkylene, C₂₋₁₆ alkenylene or phenylene group; ring Q₅ is a group of the formula:




34. A compound of claim 1 which is (R)-N-[1-(1,4-benzodioxan-2-ylmethyl)piperidin-4-ylmethyl]-5-thia-1,8b-diazaacenaphthylene-4-carboxamide, or a pharmaceutically acceptable salt thereof.

35. A compound of claim 1 which is N-[1-(3-phenylpropyl)piperidin-4-ylmethyl]-3-(5-thia-1,8b-diazaacenaphthylene-4-yl)acrylamide, or a pharmaceutically acceptable salt thereof.

36. A compound of claim 1 which is N-[4-(4-phenylpiperidin-1-yl)butan-1-yl]-5-thia-1,8b-diazaacenaphthylene-4-carboxamide, or a pharmaceutically acceptable salt thereof.

37. A compound of claim 1 which is N-[1-(3-phenylpropan-1-yl)piperidin-4-yl]-5-thia-1,8b-diazaacenaphthylene-4-carboxamide, or a pharmaceutically acceptable salt thereof.

38. A process for producing a compound of claim 5, which comprises condensing a compound of the formula:




wherein all symbols are of the same meanings as defined in claim 5, or a salt thereof with a compound of the formula:



wherein all symbols are of the same meanings as defined in claim 5, or a salt thereof.

39. A compound of the formula:



wherein R' is an optionally protected COOH, CH₂OH or CHO group; and the other symbols are of the same meanings as defined in claim 4, or a salt thereof.

40. A pharmaceutical composition which comprises a compound of claim 1.

41. A pharmaceutical composition of claim 40, which is an up-regulator of low density lipoprotein receptor.

42. A pharmaceutical composition of claim 40, which is a therapeutic agent for lowering lipids in blood.

43. A pharmaceutical composition of claim 40, which is a therapeutic agent for atherosclerosis.

44. A pharmaceutical composition of claim 40, which is an agent for lowering blood sugar.

45. A pharmaceutical composition of claim 40, which is a therapeutic agent for diabetic complications.

46. Use of a compound of claim 1 for the manufacture of a medicament for lowering lipids in blood.

47. Method for lowering lipids in blood in a mammal which comprises administering to said mammal an effective amount of a compound of claim 1.